



Certificate of Analysis

Kai Ora Honey Ltd
 56 West Road
 Awanui, Kaitaia 0483
 Attention: Blanche Morrogh
 Phone: 02108283304
 Email: blanche@kaiorahoney.co.nz

Lab Reference: 24-27259
 Submitted by:
 Date Received: 3/09/2024
 Testing Initiated: 3/09/2024
 Date Completed: 6/09/2024
 Order Number:
 Reference:

Report Comments

Samples were collected by yourselves (or your agent) and analysed as received at Analytica Laboratories (or at the subcontracted laboratories, when applicable). Samples were in acceptable condition unless otherwise noted on this report. Specific testing dates are available on request.

Results Summary

Client Sample ID		KO24FA	
Laboratory ID		24-27259-1	
Analyte	Unit	Reporting Limit	Result
MPI Manuka Classification for Honey*			
MPI Manuka Classification*			MONOFLORAL MANUKA
MPI Manuka DNA in Honey			
Manuka DNA	Cq		30.67
MPI Manuka Markers in Honey			
4-Hydroxyphenyllactic acid (4-HPLA)	mg/kg	0.80	7.5
2-Methoxybenzoic acid (2-MBA)	mg/kg	0.80	78
2'-Methoxy acetophenone (2'-MAP)	mg/kg	0.80	6.8
3-Phenyllactic acid (3-PLA)	mg/kg	20	1,700
3in1 in Honey			
Dihydroxyacetone (DHA)	mg/kg	40	2,150
Methylglyoxal (MG/MGO)	mg/kg	8	977
Non-Peroxide Activity* (NPA)	%w/v phenol eq.	1.3	22.1
Hydroxymethylfurfural (HMF)	mg/kg	1	20.0
American Foulbrood (AFB) in Honey			
AFB Cq	Cq		[ND]
AFB Amount	spores and cells per g		[ND]
Result			Not detected


 Alicia Laing, BSc.
 Technician


 Alesha Avery, BSc(Tech),
 Laboratory Technician

Method Summary

3in1	Determination of Dihydroxyacetone (DHA), Methylglyoxal (MG/MGO) and Hydroxymethylfurfural (HMF) by aqueous extraction, derivatisation, and UPLC (diode array) analysis in accordance with in-house procedures.
NPA	<p>Non-Peroxide Activity (NPA) values are not directly measured by the laboratory, but are calculated from the measured methylglyoxal concentration in the honey according to the requirements of the client. The calculation is based on published data(†) comparing the NPA and methylglyoxal concentration measured in a range of honey samples. These calculated values are not accredited by IANZ and do not imply that the honey is or is not manuka honey. NPA values less than 5 are an estimate based on extrapolation of the relationship between methylglyoxal and NPA</p> <p>(†) <i>Isolation by HPLC and characterisation of the bioactive fraction of New Zealand manuka (Leptospermum scoparium) honey. C. J. Adams, et al. Carbohydrate Research 343 (2008) 651-659. And, Corrigendum to "Isolation by HPLC and characterization of the bioactive fraction of New Zealand manuka (Leptospermum scoparium) honey" [Carbohydr. Res. 343 (2008) 651]. Carbohydrate Research 344 (2009) 2609. C. J. Adams, et al.</i></p>
MPI Manuka Markers	<p>Solvent extraction, LC-MS/MS analysis in accordance with in-house procedures.</p> <p>Analytica Laboratories Ltd., is approved by the New Zealand Ministry of Primary Industries to conduct this analysis under the Recognised Laboratory Programme (MPI Technical Paper 2017/30 Modified, RLP Method 10.05)</p>
Leptospermum scoparium DNA (PCR)	<p>Samples were analysed as received by the Laboratory for Manuka Pollen DNA by pollen DNA extraction followed by qPCR in accordance with the MPI Technical Paper 2017/31 (modified) (96 well method with magnetic bead extraction). Analytica Laboratories Ltd., is approved by the New Zealand Ministry of Primary Industries to conduct this analysis under the Recognised Laboratory Programme (RLP Method 10.04).</p> <p>The DNA component of the MPI Manuka Honey Definition requires a Cq value of less than 36 to qualify for either a monofloral or multifloral manuka honey.</p>
MPI Manuka Classification	<p>For classification as monofloral manuka, the following chemicals all need to be present and at these levels (Animal Products Notice - General Export Requirements for Bee Products, 2018):</p> <ul style="list-style-type: none">• 4-hydroxyphenyllactic acid at a level greater than or equal to 1mg/kg• 2-methoxybenzoic acid at a level greater than or equal to 1mg/kg• 2'-methoxyacetophenone at a level greater than or equal to 5mg/kg• 3-phenyllactic acid at a level greater than or equal to 400mg/kg <p>And the DNA level from manuka pollen is less than Cq 36, which is approximately 3fg/μL.</p> <p>For classification as multifloral manuka, the following chemicals all need to be present and at these levels:</p> <ul style="list-style-type: none">• 4-hydroxyphenyllactic acid at a level greater than or equal to 1mg/kg• 2-methoxybenzoic acid at a level greater than or equal to 1mg/kg• 2'-methoxyacetophenone at a level greater than or equal to 1mg/kg• 3-phenyllactic acid at a level greater than or equal to 20 mg/kg but less than 400mg/kg <p>And the DNA level from manuka pollen is less than Cq 36, which is approximately 3fg/μL.</p>
AFB	<p>Samples were analysed as received by the laboratory for AFB detection by honey DNA extraction (Magnetic Bead Method A) followed by qPCR.</p> <p>Analytica Laboratories Ltd., is approved by the New Zealand Ministry of Primary Industries to conduct this analysis under the Recognised Laboratory Programme (RLP Method 2.14).</p> <p>The LoQ (limit of quantification) is Cq 35.00 cycles (±1.5) which translates to 92 spores and/or cells per g of honey. The LoQ is not a limiting value for PCR analysis and Cq values past the LoQ cannot be considered as negative. These results are reported as Detected, as values past the LoQ can be valid in terms of presence of the target gene; however, the probability of their repeated detection is lower than 95%.</p> <p>[ND]: Not Detected.</p>